

# On a functional equation related to distributivity of fuzzy implications

Michał Baczyński, Wanda Niemyska, Tomasz Szostok

Institute of Mathematics  
University of Silesia  
Poland

May 3rd, 2013

ISCAMI, Czech Republic, May 2013

# References

- [1] M. Baczyński, W. Niemyska and T. Szostok: *On a functional equation related to distributivity of fuzzy implications*, accepted for FUZZ-IEEE 2013.

# References

- [1] M. Baczyński, W. Niemyska and T. Szostok: *On a functional equation related to distributivity of fuzzy implications*, accepted for FUZZ-IEEE 2013.
- [2] M. Baczyński and B. Jayaram: *On the distributivity of fuzzy implications over nilpotent or strict triangular conorms*, IEEE Trans. Fuzzy Syst. 17 (2009) 590–603.
- [3] W.E. Combs, J.E. Andrews: *Combinatorial rule explosion eliminated by a fuzzy rule configuration*, IEEE Trans. Fuzzy Syst. 6 (1998) 1–11.
- [4] E.P. Klement, R. Mesiar and E. Pap: *Triangular Norms*, Kluwer, Dordrecht, 2000.
- [5] M. Kuczma: *An Introduction to the Theory of Functional Equations and Inequalities. Cauchy's Equation and Jensen's Inequality*, Państwowe Wydawnictwo Naukowe (Polish Scientific Publishers) and Uniwersytet Śląski, Warszawa–Kraków–Katowice 1985.
- [6] C.H. Ling: *Representation of associative functions*, Publ. Math. Debrecen 12 (1965) 189–212.

# Content

$$f(m_1(x + y)) = m_2(f(x) + f(y))$$

# Content

$$f(m_1(x + y)) = m_2(f(x) + f(y))$$

- Some intuition what „fuzzy logic” is.

# Content

$$f(m_1(x + y)) = m_2(f(x) + f(y))$$

- Some intuition what „fuzzy logic” is.
- What do we study this equation for?

# Content

$$f(m_1(x + y)) = m_2(f(x) + f(y))$$

- Some intuition what „fuzzy logic” is.
- What do we study this equation for?
- How have we come to this equation?

# Content

$$f(m_1(x + y)) = m_2(f(x) + f(y))$$

- Some intuition what „fuzzy logic” is.
- What do we study this equation for?
- How have we come to this equation?
- The solutions of equation we have obtained.



# Content

$$f(m_1(x + y)) = m_2(f(x) + f(y))$$

- Some intuition what „fuzzy logic” is.
- What do we study this equation for?
- How have we come to this equation?
- The solutions of equation we have obtained.
- What does still remain to do?

# Classical formulas

I'm Human, so I'm Female.

# Classical formulas

I'm Human, so I'm Female.

I'm Handsome and Smart,  
so I'm Happy.

# Classical formulas

I'm Human, so I'm Female.

I'm Handsome and Smart,  
so I'm Happy.

$(\text{Handsome} \wedge \text{Smart}) \rightarrow \text{Happy} ?$

# Classical formulas

I'm Human, so I'm Female.

I'm Handsome and Smart,  
so I'm Happy.

$(\text{Handsome} \wedge \text{Smart}) \rightarrow \text{Happy} ?$

$(0.9 \wedge 0.5) \rightarrow 0.6 ?$

# Classical formulas

I'm Human, so I'm Female.

I'm Handsome and Smart,  
so I'm Happy.

$(\text{Handsome} \wedge \text{Smart}) \rightarrow \text{Happy} ?$

$(0.9 \wedge 0.5) \rightarrow 0.6 ?$

$I(T(0.9, 0.5), 0.6) = ?$

# Introduction of fuzzy operators - Implication

$$x \rightarrow y$$

$$" \rightarrow " : \{0,1\}^2 \rightarrow \{0,1\}$$

# Introduction of fuzzy operators - Implication

$$x \rightarrow y$$

$$" \rightarrow " : \{0,1\}^2 \rightarrow \{0,1\}$$

$$0 \rightarrow 0 = 1$$

$$0 \rightarrow 1 = 1$$

$$1 \rightarrow 0 = 0$$

$$1 \rightarrow 1 = 1$$



# Introduction of fuzzy operators - Implication

$$x \rightarrow y$$

$$" \rightarrow " : \{0, 1\}^2 \rightarrow \{0, 1\}$$

$$0 \rightarrow 0 = 1$$

$$0 \rightarrow 1 = 1$$

$$1 \rightarrow 0 = 0$$

$$1 \rightarrow 1 = 1$$



$$I(x, y)$$

$$I : [0, 1]^2 \rightarrow [0, 1]$$

# Introduction of fuzzy operators - Implication

$$x \rightarrow y$$

$$" \rightarrow " : \{0, 1\}^2 \rightarrow \{0, 1\}$$

$$0 \rightarrow 0 = 1$$

$$0 \rightarrow 1 = 1$$

$$1 \rightarrow 0 = 0$$

$$1 \rightarrow 1 = 1$$



$$I(x, y)$$

$$I : [0, 1]^2 \rightarrow [0, 1]$$

$$I(0, 0) = 1$$

$$I(0, 1) = 1$$

$$I(1, 0) = 0$$

$$I(1, 1) = 1$$

# Introduction of fuzzy operators - Implication

$$x \rightarrow y$$

$$" \rightarrow " : \{0, 1\}^2 \rightarrow \{0, 1\}$$

$$0 \rightarrow 0 = 1$$

$$0 \rightarrow 1 = 1$$

$$1 \rightarrow 0 = 0$$

$$1 \rightarrow 1 = 1$$



$$I(x, y)$$

$$I : [0, 1]^2 \rightarrow [0, 1]$$

$$I(0, 0) = 1$$

$$I(0, 1) = 1$$

$$I(1, 0) = 0$$

$$I(1, 1) = 1$$

## Definition 1

A function  $I : [0, 1]^2 \rightarrow [0, 1]$  is called a **FUZZY IMPLICATION** if it satisfies, for all  $x, y \in [0, 1]$ , the following conditions

- 1  $I(\cdot, y)$  is decreasing,
- 2  $I(x, \cdot)$  is increasing,
- 3  $I(0, 0) = 1$ ,
- 4  $I(1, 1) = 1$ ,
- 5  $I(1, 0) = 0$ .

# Introduction of fuzzy operators - Conjunction

$$x \wedge y$$

$$" \wedge ": \{0, 1\}^2 \rightarrow \{0, 1\}$$

$$0 \wedge 0 = 0$$

$$0 \wedge 1 = 0$$

$$1 \wedge 0 = 0$$

$$1 \wedge 1 = 1$$



$$T(x, y)$$

$$T: [0, 1]^2 \rightarrow [0, 1]$$

$$T(0, 0) = 0$$

$$T(0, 1) = 0$$

$$T(1, 0) = 0$$

$$T(1, 1) = 1$$

## Definition 2

A function  $T: [0, 1]^2 \rightarrow [0, 1]$  is called a **TRIANGULAR NORM (t-norm)** if it satisfies, for all  $x, y, z \in [0, 1]$ , the following conditions

- 1  $T(x, y) = T(y, x)$ ,
- 2  $T(x, T(y, z)) = T(T(x, y), z)$ ,
- 3  $T(x, \cdot)$  is decreasing,
- 4  $T(x, 1) = x$ .

# Introduction of fuzzy operators - Disjunction

$$x \vee y$$

$$" \vee ": \{0,1\}^2 \rightarrow \{0,1\}$$

$$0 \vee 0 = 0$$

$$0 \vee 1 = 1$$

$$1 \vee 0 = 1$$

$$1 \vee 1 = 1$$



$$S(x, y)$$

$$S: [0,1]^2 \rightarrow [0,1]$$

$$S(0,0) = 0$$

$$S(0,1) = 1$$

$$S(1,0) = 1$$


$$S(1,1) = 1$$

## Definition 3


A function  $S: [0,1]^2 \rightarrow [0,1]$  is called a **TRIANGULAR CONORM** (*t-conorm*) if it satisfies, for all  $x, y, z \in [0,1]$ , the following conditions

- 1  $S(x, y) = S(y, x)$ ,
- 2  $S(x, S(y, z)) = S(S(x, y), z)$ ,
- 3  $S(x, \cdot)$  is increasing,
- 4  $S(x, 0) = x$ .

# Fuzzy thinking



# Fuzzy thinking




# Fuzzy thinking






# Fuzzy thinking



# Fuzzy thinking



# Compositional Rule of Inference (CRI): Laundry

# Compositional Rule of Inference (CRI): Laundry

- **D** - the laundry is Dirty;

# Compositional Rule of Inference (CRI): Laundry

- **D** - the laundry is Dirty;
- **VD** - the laundry is Very Dirty;

# Compositional Rule of Inference (CRI): Laundry

- **D** - the laundry is Dirty;
- **VD** - the laundry is Very Dirty;
- **S** - the laundry is Small;

# Compositional Rule of Inference (CRI): Laundry

- **D** - the laundry is Dirty;
- **VD** - the laundry is Very Dirty;
- **S** - the laundry is Small;
- **B** - the laundry is Big;

# Compositional Rule of Inference (CRI): Laundry

- **D** - the laundry is Dirty;
- **VD** - the laundry is Very Dirty;
- **S** - the laundry is Small;
- **B** - the laundry is Big;
- **T<sub>>x</sub>** - at least  $x$  minutes of laundering;





# Compositional Rule of Inference (CRI): Laundry

- **D** - the laundry is Dirty;
- **VD** - the laundry is Very Dirty;
- **S** - the laundry is Small;
- **B** - the laundry is Big;
- **T**<sub>>x</sub> - at least x minutes of laundering;



$$(D \wedge S) \rightarrow T_{>45}$$

$$(VD \wedge S) \rightarrow T_{>60}$$

$$(D \wedge B) \rightarrow T_{>70}$$

$$(VD \wedge B) \rightarrow T_{>90}$$

# Compositional Rule of Inference (CRI): Laundry

- **D** - the laundry is Dirty;
- **VD** - the laundry is Very Dirty;
- **S** - the laundry is Small;
- **B** - the laundry is Big;
- **T<sub>>x</sub>** - at least x minutes of laundering;



$$(D \wedge S) \rightarrow T_{>45}$$

$$(VD \wedge S) \rightarrow T_{>60}$$

$$(D \wedge B) \rightarrow T_{>70}$$

$$(VD \wedge B) \rightarrow T_{>90}$$

$$(D \rightarrow T_{>45}) \vee (S \rightarrow T_{>45})$$

$$(VD \rightarrow T_{>60}) \vee (S \rightarrow T_{>60})$$

$$(D \rightarrow T_{>70}) \vee (B \rightarrow T_{>70})$$

$$(VD \rightarrow T_{>90}) \vee (B \rightarrow T_{>90})$$

$$(x \wedge y) \rightarrow z \equiv (x \rightarrow z) \vee (y \rightarrow z) \quad (T1)$$

# Compositional Rule of Inference (CRI): Laundry

$$x \rightarrow (y \vee z) \equiv (x \rightarrow y) \vee (x \rightarrow z) \quad (T2)$$

$$D \rightarrow T_{>45}, \quad S \rightarrow T_{>45},$$

$$VD \rightarrow T_{>60}, \quad S \rightarrow T_{>60},$$

$$D \rightarrow T_{>70}, \quad B \rightarrow T_{>70},$$

$$VD \rightarrow T_{>90}, \quad B \rightarrow T_{>90}$$

# Compositional Rule of Inference (CRI): Laundry

$$x \rightarrow (y \vee z) \equiv (x \rightarrow y) \vee (x \rightarrow z) \quad (T2)$$

$$\begin{array}{lll} D \rightarrow T_{>45}, & S \rightarrow T_{>45}, & D \rightarrow (T_{>45} \vee T_{>70}) \\ VD \rightarrow T_{>60}, & S \rightarrow T_{>60}, & \\ D \rightarrow T_{>70}, & B \rightarrow T_{>70}, & \\ VD \rightarrow T_{>90}, & B \rightarrow T_{>90} & \end{array}$$

# Compositional Rule of Inference (CRI): Laundry

$$x \rightarrow (y \vee z) \equiv (x \rightarrow y) \vee (x \rightarrow z) \quad (T2)$$

$D \rightarrow T_{>45},$	$S \rightarrow T_{>45},$	$D \rightarrow (T_{>45} \vee T_{>70})$
$VD \rightarrow T_{>60},$	$S \rightarrow T_{>60},$	$VD \rightarrow (T_{>60} \vee T_{>90})$
$D \rightarrow T_{>70},$	$B \rightarrow T_{>70},$	$S \rightarrow (T_{>45} \vee T_{>60})$
$VD \rightarrow T_{>90},$	$B \rightarrow T_{>90}$	$B \rightarrow (T_{>70} \vee T_{>90})$

# Compositional Rule of Inference (CRI): Laundry

$$x \rightarrow (y \vee z) \equiv (x \rightarrow y) \vee (x \rightarrow z) \quad (T2)$$


$D \rightarrow T_{>45},$	$S \rightarrow T_{>45},$	$D \rightarrow (T_{>45} \vee T_{>70})$	$D \rightarrow T_{>45}$
$VD \rightarrow T_{>60},$	$S \rightarrow T_{>60},$	$VD \rightarrow (T_{>60} \vee T_{>90})$	$VD \rightarrow T_{>60}$
$D \rightarrow T_{>70},$	$B \rightarrow T_{>70},$	$S \rightarrow (T_{>45} \vee T_{>60})$	$S \rightarrow T_{>45}$
$VD \rightarrow T_{>90},$	$B \rightarrow T_{>90}$	$B \rightarrow (T_{>70} \vee T_{>90})$	$B \rightarrow T_{>70}$

# Compositional Rule of Inference (CRI): Laundry

IF the laundry is Big, THEN ...

# Compositional Rule of Inference (CRI): Laundry

IF the laundry is Big, THEN ...





# Compositional Rule of Inference (CRI): Laundry

IF the laundry is Big, THEN ....



# Compositional Rule of Inference (CRI): Laundry

IF the laundry is Big, THEN ....



$$\text{Big}(\text{Laundry1}) = 1$$



$$\text{Big}(\text{Laundry2}) = 0$$



$$\text{Big}(\text{Laundry3}) = 0.27$$



$$\text{Big}(\text{Laundry4}) = 0.62$$

# Compositional Rule of Inference (CRI) - Laundry

The general schema of multiconditional approximate reasoning has the form:

*Rule 1:* IF  $\tilde{x}$  is  $A_1$ , THEN  $\tilde{y}$  is  $B_1$

*Rule 2:* IF  $\tilde{x}$  is  $A_2$ , THEN  $\tilde{y}$  is  $B_2$

.....

*Rule n:* IF  $\tilde{x}$  is  $A_n$ , THEN  $\tilde{y}$  is  $B_n$

*Fact:*  $\tilde{x}$  is  $A$

# Compositional Rule of Inference (CRI) - Laundry

The general schema of multiconditional approximate reasoning has the form:

*Rule 1:* IF  $\tilde{x}$  is  $A_1$ , THEN  $\tilde{y}$  is  $B_1$

*Rule 2:* IF  $\tilde{x}$  is  $A_2$ , THEN  $\tilde{y}$  is  $B_2$

.....

*Rule n:* IF  $\tilde{x}$  is  $A_n$ , THEN  $\tilde{y}$  is  $B_n$

*Fact:*  $\tilde{x}$  is  $A$

*Conclusion:*  $\tilde{y}$  is  $B$

$$B(y) = \sup_{j \in N_n} \sup_{x \in X} T(A(x), I(A_j(x), B_j(y)))$$

# Compositional Rule of Inference (CRI) - Laundry

The general schema of multiconditional approximate reasoning has the form:

*Rule 1:* IF  $\tilde{x}$  is  $A_1$ , THEN  $\tilde{y}$  is  $B_1$

*Rule 2:* IF  $\tilde{x}$  is  $A_2$ , THEN  $\tilde{y}$  is  $B_2$

.....

*Rule n:* IF  $\tilde{x}$  is  $A_n$ , THEN  $\tilde{y}$  is  $B_n$

*Fact:*  $\tilde{x}$  is  $A$

*Conclusion:*  $\tilde{y}$  is  $B$

$$B(y) = \sup_{j \in N_n} \sup_{x \in X} T(A(x), I(A_j(x), B_j(y)))$$



IF  $\tilde{x}$  is **D**, THEN  $\tilde{y}$  is  $\mathbf{T}_{>45}$

IF  $\tilde{x}$  is **VD**, THEN  $\tilde{y}$  is  $\mathbf{T}_{>60}$

IF  $\tilde{x}$  is **S**, THEN  $\tilde{y}$  is  $\mathbf{T}_{>45}$

IF  $\tilde{x}$  is **B**, THEN  $\tilde{y}$  is  $\mathbf{T}_{>70}$

# Derivation of the functional equation

$$(x \wedge y) \rightarrow z \equiv (x \rightarrow z) \vee (y \rightarrow z) \quad (T1)$$

$$x \rightarrow (y \vee z) \equiv (x \rightarrow y) \vee (x \rightarrow z) \quad (T2)$$

# Derivation of the functional equation

$$(x \wedge y) \rightarrow z \equiv (x \rightarrow z) \vee (y \rightarrow z) \quad (T1)$$

$$x \rightarrow (y \vee z) \equiv (x \rightarrow y) \vee (x \rightarrow z) \quad (T2)$$



$$I(T(x, y), z) = S(I(x, z), I(y, z)) \quad (D1)$$

$$I(x, S(y, z)) = S(I(x, y), I(x, z)) \quad (D2)$$

# Derivation of the functional equation

$$(x \wedge y) \rightarrow z \equiv (x \rightarrow z) \vee (y \rightarrow z) \quad (T1)$$

$$x \rightarrow (y \vee z) \equiv (x \rightarrow y) \vee (x \rightarrow z) \quad (T2)$$



$$I(T(x, y), z) = S(I(x, z), I(y, z)) \quad (D1)$$

$$I(x, S(y, z)) = S(I(x, y), I(x, z)) \quad (D2)$$



# Derivation of the functional equation

$$x \rightarrow (y \vee z) \equiv (x \rightarrow y) \vee (x \rightarrow z) \quad (T2)$$

TAUTOLOGY

# Derivation of the functional equation

$$x \rightarrow (y \vee z) \equiv (x \rightarrow y) \vee (x \rightarrow z) \quad (T2)$$

## TAUTOLOGY

x	y	z	$x \rightarrow (y \vee z)$		$(x \rightarrow y) \vee (x \rightarrow z)$			$L \equiv R$
			$y \vee z$	$L$	$x \rightarrow y$	$x \rightarrow z$	$R$	
0	0	0	0	1	1	1	1	1
0	0	1	1	1	1	1	1	1
0	1	0	1	1	1	1	1	1
1	0	0	0	0	0	0	0	1
0	1	1	1	1	1	1	1	1
1	0	1	1	1	0	1	1	1
1	1	0	1	1	1	0	1	1
1	1	1	1	1	1	1	1	1

# Derivation of the functional equation

$$I(x, S(y, z)) = S(I(x, y), I(x, z)) \quad (D2)$$

# Derivation of the functional equation

$$I(x, S(y, z)) = S(I(x, y), I(x, z)) \quad (D2)$$

## TAUTOLOGY?

**The question:** When the above equation is true? For which  $S$  and  $I$ ?

# Derivation of the functional equation

$$I(x, S(y, z)) = S(I(x, y), I(x, z)) \quad (D2)$$

## TAUTOLOGY?

**The question:** When the above equation is true? For which  $S$  and  $I$ ?

$$I(x, S_1(y, z)) = S_2(I(x, y), I(x, z))$$

# Derivation of the functional equation

$$I(x, S(y, z)) = S(I(x, y), I(x, z)) \quad (D2)$$

## TAUTOLOGY?

**The question:** When the above equation is true? For which  $S$  and  $I$ ?

$$I(x, S_1(y, z)) = S_2(I(x, y), I(x, z))$$

### Lemma 1

*When  $S$  is continuous and Archimedean triangular conorm, then  $S$  is of the form*

$$S(x, y) = s^{-1}(\min(s(x) + s(y), s(1))), \quad x, y \in [0, 1]$$

*where  $s : [0, 1] \rightarrow [0, \infty]$  is a continuous, strictly increasing function with  $s(0) = 0$ .*

# Derivation of the functional equation

$$I(x, S_1(y, z)) = S_2(I(x, y), I(x, z))$$

# Derivation of the functional equation

$$I(x, S_1(y, z)) = S_2(I(x, y), I(x, z))$$



***Lemma 1*** and some simple calculations





# Derivation of the functional equation

$$I(x, S_1(y, z)) = S_2(I(x, y), I(x, z))$$



**Lemma 1** and some simple calculations



$$f(\min(u + v, r_1)) = \min(f(u) + f(v), r_2),$$

where  $u, v \in [0, r_1]$ ,  $r_1 := s_1(1)$ ,  $r_2 := s_2(1) \in (0, \infty)$

# Derivation of the functional equation

$$I(x, S_1(y, z)) = S_2(I(x, y), I(x, z))$$



**Lemma 1** and some simple calculations



$$f(\min(u + v, r_1)) = \min(f(u) + f(v), r_2),$$

where  $u, v \in [0, r_1]$ ,  $r_1 := s_1(1)$ ,  $r_2 := s_2(1) \in (0, \infty)$

- For  $S_1, S_2$  both nilpotent or both strict  $t$ -conorms we know the form of implication  $I$ ,
- For a  $R$ -implication  $I$  generated from a strict  $t$ -norm  $T$  equation holds if and only if  $t$ -conorms  $S_1 = S_2$  are  $\Phi$ -conjugate with the Łukasiewicz  $t$ -conorm for some increasing bijection  $\Phi$ , which is a multiplicative generator of the strict  $t$ -norm  $T$ .

# Derivation of the functional equation

$$f(\min(u + v, r_1)) = \min(f(u) + f(v), r_2),$$

where  $u, v \in [0, r_1]$ ,  $r_1 := s_1(1), r_2 := s_2(1) \in (0, \infty)$

# Derivation of the functional equation

$$f(\min(u + v, r_1)) = \min(f(u) + f(v), r_2),$$

where  $u, v \in [0, r_1]$ ,  $r_1 := s_1(1)$ ,  $r_2 := s_2(1) \in (0, \infty)$

↓

$$f(m_1(x + y)) = m_2(f(x) + f(y)) \quad (1),$$

where for fixed  $r_1, r_2 \in (0, \infty)$  functions  
 $m_1 : [0, 2r_1] \rightarrow [0, r_1]$ ,  $m_2 : [0, 2r_2] \rightarrow [0, r_2]$  and  $f : [0, r_1] \rightarrow [0, r_2]$ .

# $m_2$ - injective

## Theorem 1

*Let  $r_1, r_2 \in (0, \infty)$  be some numbers and let  $m_1: [0, 2r_1] \rightarrow [0, r_1]$ ,  $m_2: [0, 2r_2] \rightarrow [0, r_2]$ ,  $f: [0, r_1] \rightarrow [0, r_2]$  be given functions. Further, let  $m_2$  be injective. Then the following sentences are equivalent:*

# $m_2$ - injective

## Theorem 1

*Let  $r_1, r_2 \in (0, \infty)$  be some numbers and let  $m_1: [0, 2r_1] \rightarrow [0, r_1]$ ,  $m_2: [0, 2r_2] \rightarrow [0, r_2]$ ,  $f: [0, r_1] \rightarrow [0, r_2]$  be given functions. Further, let  $m_2$  be injective. Then the following sentences are equivalent:*

- 1** *The triple of functions  $m_1, m_2, f$  satisfies the equation (1).*

# $m_2$ - injective

## Theorem 1

*Let  $r_1, r_2 \in (0, \infty)$  be some numbers and let  $m_1: [0, 2r_1] \rightarrow [0, r_1]$ ,  $m_2: [0, 2r_2] \rightarrow [0, r_2]$ ,  $f: [0, r_1] \rightarrow [0, r_2]$  be given functions. Further, let  $m_2$  be injective. Then the following sentences are equivalent:*

- 1** *The triple of functions  $m_1, m_2, f$  satisfies the equation (1).*
- 2**  *$f(x) = ax + b$  for some  $a, b \in \mathbb{R}$ .*

# $m_2$ - injective

## Theorem 1

Let  $r_1, r_2 \in (0, \infty)$  be some numbers and let  $m_1: [0, 2r_1] \rightarrow [0, r_1]$ ,  $m_2: [0, 2r_2] \rightarrow [0, r_2]$ ,  $f: [0, r_1] \rightarrow [0, r_2]$  be given functions. Further, let  $m_2$  be injective. Then the following sentences are equivalent:

- 1 The triple of functions  $m_1, m_2, f$  satisfies the equation (1).
- 2  $f(x) = ax + b$  for some  $a, b \in \mathbb{R}$ .

More precisely:

*Either  $f = b$  for some  $b \in [0, r_2]$  and  $m_2(2b) = b$ , or  $f(x) = ax + b$  for some  $a, b \in \mathbb{R}$ ,  $a \neq 0$  such that*

$$ax + b \in [0, r_2], \quad \text{for all } x \in [0, r_1] \quad (1)$$

and

$$m_1(x) = \frac{m_2(ax + 2b) - b}{a}. \quad (2)$$



# $m_2$ - NOT injective

## Theorem 2

*Let  $r_1, r_2 \in (0, \infty)$  be some numbers and let functions  $m_1: [0, 2r_1] \rightarrow [0, r_1]$ ,  $m_2: [0, 2r_2] \rightarrow [0, r_2]$  be continuous and strictly increasing on some intervals  $[0, x_1]$ ,  $[0, x_2]$ , respectively, and then be equal to  $r_1$ ,  $r_2$ , respectively, where  $x_1 \leq r_1$  and  $x_2 \leq r_2$ . Further, let  $m_1, m_2$  satisfy*

$$m_1(0) = 0, \quad 2m_1(x) > x, \quad x \in (0, 2r_1) \quad (3)$$

*and*

$$m_2(0) = 0, \quad 2m_2(x) > x, \quad x \in (0, 2r_2). \quad (4)$$

*. Finally let  $f$  be a function  $f: [0, r_1] \rightarrow [0, r_2]$ .*

# $m_2$ - NOT injective

## Theorem 2

*Let  $f, m_1, m_2$  satisfy the above assumptions.*

*If that triple of functions satisfies equation (1), then we have one of the following possibilities:*

# $m_2$ - NOT injective

## Theorem 2

*Let  $f, m_1, m_2$  satisfy the above assumptions.*

*If that triple of functions satisfies equation (1), then we have one of the following possibilities:*

- 1  $f = r_2$  and  $m_1, m_2$  may be any functions;

# $m_2$ - NOT injective

## Theorem 2

*Let  $f, m_1, m_2$  satisfy the above assumptions.*

*If that triple of functions satisfies equation (1), then we have one of the following possibilities:*

- 1**  $f = r_2$  and  $m_1, m_2$  may be any functions;
- 2**  $f = 0$  and  $m_1, m_2$  may be any functions;

# $m_2$ - NOT injective

## Theorem 2

*Let  $f, m_1, m_2$  satisfy the above assumptions.*

*If that triple of functions satisfies equation (1), then we have one of the following possibilities:*

- 1**  $f = r_2$  and  $m_1, m_2$  may be any functions;
- 2**  $f = 0$  and  $m_1, m_2$  may be any functions;
- 3**  $f(0) = 0, f(x) > x_2$  for  $x > 0, f(r_1) = r_2$ , and  $m_1, m_2$  may be any functions;

# $m_2$ - NOT injective

## Theorem 2

*Let  $f, m_1, m_2$  satisfy the above assumptions.*

*If that triple of functions satisfies equation (1), then we have one of the following possibilities:*

- 1**  $f = r_2$  and  $m_1, m_2$  may be any functions;
- 2**  $f = 0$  and  $m_1, m_2$  may be any functions;
- 3**  $f(0) = 0, f(x) > x_2$  for  $x > 0, f(r_1) = r_2$ , and  $m_1, m_2$  may be any functions;
- 4** there exists  $x_0 \in (0, r_1]$  such that  $f(x) \geq x_2$  for  $x \geq x_0, f(x) = r_2$  for  $x \in [m_1(x_0), r_1]$  and  $f(x) = \frac{x_2}{x_0}x$  for  $x < x_0$ . Moreover in this case

$$m_1(x) = \frac{x_0 m_2\left(\frac{x_2}{x_0}x\right)}{x_2} \quad \text{for } x < y_0, \text{ such that } m_1(y_0) = x_0.$$

# $m_2$ - NOT injective

## Theorem 2

*Let  $f, m_1, m_2$  satisfy the above assumptions.*

*If that triple of functions satisfies equation (1), then we have one of the following possibilities:*

- 1**  $f = r_2$  and  $m_1, m_2$  may be any functions;
- 2**  $f = 0$  and  $m_1, m_2$  may be any functions;
- 3**  $f(0) = 0, f(x) > x_2$  for  $x > 0, f(r_1) = r_2$ , and  $m_1, m_2$  may be any functions;
- 4** there exists  $x_0 \in (0, r_1]$  such that  $f(x) \geq x_2$  for  $x \geq x_0, f(x) = r_2$  for  $x \in [m_1(x_0), r_1]$  and  $f(x) = \frac{x_2}{x_0}x$  for  $x < x_0$ . Moreover in this case

$$m_1(x) = \frac{x_0 m_2\left(\frac{x_2}{x_0}x\right)}{x_2} \quad \text{for } x < y_0, \text{ such that } m_1(y_0) = x_0.$$

*Conversely, if we add to the case (4) an assumption that  $y_0 = x_0$  or  $f(m_1(x)) = m_2(f(x))$  for  $x \in [y_0, x_0)$ , then each of the triples of functions described above satisfies the equation (1).*

# $m_2$ - NOT injective

## Remark 1

*We've showed that an additional assumption in the converse to Theorem 2 ( $y_0 = x_0$  or  $f(m_1(x)) = m_2(f(x))$  for  $x \in [y_0, x_0]$ ) is necessary.*



# Example 1

## Example 1

Let us fix arbitrarily  $r_1, r_2 > 0$  and  $\alpha \geq 1$ .

- $m_1(x) = \min(\alpha x, r_1)$  for  $x \in [0, 2r_1]$
- $m_2(x) = \min(\alpha x, r_2)$  for  $x \in [0, 2r_2]$ .

In this case we obtain the following equation

$$f(\min(\alpha(x+y), r_1)) = \min(\alpha(f(x) + f(y)), r_2).$$

# Example 1

## Example 1




Let us fix arbitrarily  $r_1, r_2 > 0$  and  $\alpha \geq 1$ .

- $m_1(x) = \min(\alpha x, r_1)$  for  $x \in [0, 2r_1]$
- $m_2(x) = \min(\alpha x, r_2)$  for  $x \in [0, 2r_2]$ .

In this case we obtain the following equation

$$f(\min(\alpha(x+y), r_1)) = \min(\alpha(f(x) + f(y)), r_2).$$

From Theorem 2 we obtain that the only nontrivial continuous solution is  $f(x) = \min(kx, r_2)$ , where  $k = \frac{r_2}{\alpha x_0}$ .



## Example 2

### Example 2

Let us fix arbitrarily  $r_1, r_2 > 0$ .

- $m_1(x) = \min(\sqrt{r_1 x}, r_1)$  for  $x \in [0, 2r_1]$
- $m_2(x) = \min(\sqrt{r_2 x}, r_2)$  for  $x \in [0, 2r_2]$ .

In this case we obtain the following equation

$$f(\min(\sqrt{r_1(x+y)}, r_1)) = \min(\sqrt{r_2(f(x) + f(y))}, r_2) \quad (3)$$

# Example 2

## Example 2

Let us fix arbitrarily  $r_1, r_2 > 0$ .




- $m_1(x) = \min(\sqrt{r_1 x}, r_1)$  for  $x \in [0, 2r_1]$
- $m_2(x) = \min(\sqrt{r_2 x}, r_2)$  for  $x \in [0, 2r_2]$ .

In this case we obtain the following equation

$$f(\min(\sqrt{r_1(x+y)}, r_1)) = \min(\sqrt{r_2(f(x) + f(y))}, r_2) \quad (3)$$

From Theorem 2 we obtain that the only nontrivial continuous solution is

$$f(x) = \frac{r_2}{r_1} x.$$



# Conclusion

We have discussed some solutions of one functional equation which generalizes an equation connected with solutions of the distributivity equation of fuzzy implications functions over some classes of triangular conorms:



# Conclusion

We have discussed some solutions of one functional equation which generalizes an equation connected with solutions of the distributivity equation of fuzzy implications functions over some classes of triangular conorms:

- for  $m_2$ , which is injective, we gave complete characterization of solutions of the equation (1)



# Conclusion

We have discussed some solutions of one functional equation which generalizes an equation connected with solutions of the distributivity equation of fuzzy implications functions over some classes of triangular conorms:

- for  $m_2$ , which is injective, we gave complete characterization of solutions of the equation (1)
- for  $m_2$ , which is NOT injective, we gave partial characterization of solutions of the equation (1)



# Conclusion

We have discussed some solutions of one functional equation which generalizes an equation connected with solutions of the distributivity equation of fuzzy implications functions over some classes of triangular conorms:

- for  $m_2$ , which is injective, we gave complete characterization of solutions of the equation (1)
- for  $m_2$ , which is NOT injective, we gave partial characterization of solutions of the equation (1)

We believe that obtained results can be used in future for instance in:





# Conclusion

We have discussed some solutions of one functional equation which generalizes an equation connected with solutions of the distributivity equation of fuzzy implications functions over some classes of triangular conorms:

- for  $m_2$ , which is injective, we gave complete characterization of solutions of the equation (1)
- for  $m_2$ , which is NOT injective, we gave partial characterization of solutions of the equation (1)

We believe that obtained results can be used in future for instance in:

- fuzzy control,



# Conclusion

We have discussed some solutions of one functional equation which generalizes an equation connected with solutions of the distributivity equation of fuzzy implications functions over some classes of triangular conorms:

- for  $m_2$ , which is injective, we gave complete characterization of solutions of the equation (1)
- for  $m_2$ , which is NOT injective, we gave partial characterization of solutions of the equation (1)

We believe that obtained results can be used in future for instance in:

- fuzzy control,
- fuzzy mathematical morphology,



# Conclusion


We have discussed some solutions of one functional equation which generalizes an equation connected with solutions of the distributivity equation of fuzzy implications functions over some classes of triangular conorms:

- for  $m_2$ , which is injective, we gave complete characterization of solutions of the equation (1)
- for  $m_2$ , which is NOT injective, we gave partial characterization of solutions of the equation (1)

We believe that obtained results can be used in future for instance in:

- fuzzy control,
- fuzzy mathematical morphology,
- aggregations functions.





THANK  
YOU  
FOR YOUR  
ATTENTION!