

Upper bound on Abelian lifts of complete graphs

Dávid Mesežnikov

Slovak University of Technology

12. 5. 2012

The degree-diameter problem

The degree-diameter problem

The problem:

The degree-diameter problem

The problem:

Find the largest order $n(d, k)$ of a graph of maximum degree d and diameter k (the (d, k) -graphs) and characterize the extremal graphs.

The degree-diameter problem

The problem:

Find the largest order $n(d, k)$ of a graph of maximum degree d and diameter k (the (d, k) -graphs) and characterize the extremal graphs.

Two mainstreams of research:

The degree-diameter problem

The problem:

Find the largest order $n(d, k)$ of a graph of maximum degree d and diameter k (the (d, k) -graphs) and characterize the extremal graphs.

Two mainstreams of research:

- **Proofs of non-existence** of (d, k) -graphs of order 'close' to the **Moore bound**

$$M(d, k) = 1 + d + d(d - 1) + \dots + d(d - 1)^{k-1}$$

(except sporadic examples).

The degree-diameter problem

The problem:

Find the largest order $n(d, k)$ of a graph of maximum degree d and diameter k (the (d, k) -graphs) and characterize the extremal graphs.

Two mainstreams of research:

- **Proofs of non-existence** of (d, k) -graphs of order 'close' to the **Moore bound**

$$M(d, k) = 1 + d + d(d - 1) + \dots + d(d - 1)^{k-1}$$

(except sporadic examples).

- **Construction methods** for lower bounds on $n(d, k)$.

Non-existence - general information

Non-existence - general information

For $d \geq 3$, $k \geq 2$ we have $n(d, k) = M(d, k)$ only if (d, k) is equal to

Non-existence - general information

For $d \geq 3$, $k \geq 2$ we have $n(d, k) = M(d, k)$ only if (d, k) is equal to

- $(3, 2)$ – the Petersen graph, or

Non-existence - general information

For $d \geq 3$, $k \geq 2$ we have $n(d, k) = M(d, k)$ only if (d, k) is equal to

- $(3, 2)$ – the Petersen graph, or
- $(7, 2)$ – the Hoffman-Singleton graph, or possibly

Non-existence - general information

For $d \geq 3$, $k \geq 2$ we have $n(d, k) = M(d, k)$ only if (d, k) is equal to

- $(3, 2)$ – the Petersen graph, or
- $(7, 2)$ – the Hoffman-Singleton graph, or possibly
- $(57, 2)$ – ???

Non-existence - general information

For $d \geq 3$, $k \geq 2$ we have $n(d, k) = M(d, k)$ only if (d, k) is equal to

- $(3, 2)$ – the Petersen graph, or
- $(7, 2)$ – the Hoffman-Singleton graph, or possibly
- $(57, 2)$ – ???

For other d, k we have $n(d, k) \leq M(d, k) - 2$ and $n(3, k) \leq M(3, k) - 4$.

Construction method - Voltage assignments

Construction method - Voltage assignments

G - an undirected graph; Γ - an arbitrary finite group.

Construction method - Voltage assignments

G - an undirected graph; Γ - an arbitrary finite group.

$D(G)$ - set of all arcs of G

Construction method - Voltage assignments

G - an undirected graph; Γ - an arbitrary finite group.

$D(G)$ - set of all arcs of G

A mapping $\alpha : D(G) \rightarrow \Gamma$ is a **voltage assignment** if

$$\alpha(e^{-1}) = (\alpha(e))^{-1} \text{ for each } e \in D(G).$$

Construction method - Voltage assignments

G - an undirected graph; Γ - an arbitrary finite group.

$D(G)$ - set of all arcs of G

A mapping $\alpha : D(G) \rightarrow \Gamma$ is a **voltage assignment** if

$$\alpha(e^{-1}) = (\alpha(e))^{-1} \text{ for each } e \in D(G).$$

A graph G^α with $V(G^\alpha) = V(G) \times \Gamma$ and $D(G^\alpha) = D(G) \times \Gamma$ is called a **lift** of G .

Construction method - Voltage assignments

G - an undirected graph; Γ - an arbitrary finite group.

$D(G)$ - set of all arcs of G

A mapping $\alpha : D(G) \rightarrow \Gamma$ is a **voltage assignment** if

$$\alpha(e^{-1}) = (\alpha(e))^{-1} \text{ for each } e \in D(G).$$

A graph G^α with $V(G^\alpha) = V(G) \times \Gamma$ and $D(G^\alpha) = D(G) \times \Gamma$ is called a **lift** of G .

A vertex (u, g) of G^α is connected with (v, h) iff $g\alpha(e) = h$ and e is an edge from u to v in the base graph.

Construction method - Voltage assignments

G - an undirected graph; Γ - an arbitrary finite group.

$D(G)$ - set of all arcs of G

A mapping $\alpha : D(G) \rightarrow \Gamma$ is a **voltage assignment** if


$$\alpha(e^{-1}) = (\alpha(e))^{-1} \text{ for each } e \in D(G).$$

A graph G^α with $V(G^\alpha) = V(G) \times \Gamma$ and $D(G^\alpha) = D(G) \times \Gamma$ is called a **lift** of G .

A vertex (u, g) of G^α is connected with (v, h) iff $g\alpha(e) = h$ and e is an edge from u to v in the base graph.

The set $\{(u, g), g \in \Gamma\}$ forms a fiber above u .

Example of a lift:



McKay-Miller-Širáň graphs

McKay-Miller-Širáň graphs

The largest currently known vertex-transitive non-Cayley graphs of diameter two and given degree are the **McKay-Miller-Širáň graphs** $\text{MMS}(q)$.

McKay-Miller-Širáň graphs

The largest currently known vertex-transitive non-Cayley graphs of diameter two and given degree are the **McKay-Miller-Širáň graphs** $\text{MMS}(q)$.

These graphs were constructed as lifts of dipoles in Abelian groups. They are defined for any prime power q such that $q \equiv 1 \pmod{4}$, and have order $2q^2$, degree $d = (3q - 1)/2$, and diameter 2.

McKay-Miller-Širáň graphs

The largest currently known vertex-transitive non-Cayley graphs of diameter two and given degree are the McKay-Miller-Širáň graphs $\text{MMS}(q)$.

These graphs were constructed as lifts of dipoles in Abelian groups. They are defined for any prime power q such that $q \equiv 1 \pmod{4}$, and have order $2q^2$, degree $d = (3q - 1)/2$, and diameter 2.

In terms of degree, the order of $\text{MMS}(q)$ is $\frac{8}{9}(d + \frac{1}{2})^2$.

McKay-Miller-Širáň graphs

The largest currently known vertex-transitive non-Cayley graphs of diameter two and given degree are the **McKay-Miller-Širáň graphs** $MMS(q)$.

These graphs were constructed as lifts of dipoles in Abelian groups. They are defined for any prime power q such that $q \equiv 1 \pmod{4}$, and have order $2q^2$, degree $d = (3q - 1)/2$, and diameter 2.

In terms of degree, the order of $MMS(q)$ is $\frac{8}{9}(d + \frac{1}{2})^2$.

An interesting aside: $MMS(5)$ is the Hoffman-Singleton graph.

McKay-Miller-Širáň graphs

The largest currently known vertex-transitive non-Cayley graphs of diameter two and given degree are the McKay-Miller-Širáň graphs $\text{MMS}(q)$.

These graphs were constructed as lifts of dipoles in Abelian groups. They are defined for any prime power q such that $q \equiv 1 \pmod{4}$, and have order $2q^2$, degree $d = (3q - 1)/2$, and diameter 2.

In terms of degree, the order of $\text{MMS}(q)$ is $\frac{8}{9}(d + \frac{1}{2})^2$.

An interesting aside: $\text{MMS}(5)$ is the Hoffman-Singleton graph.

Lifts of dipoles using Abelian groups

Lifts of dipoles using Abelian groups

A dipole lift construction of MMS graphs motivates the study of largest possible graphs of diameter 2 obtained as lifts of dipoles using voltages in Abelian groups.

Lifts of dipoles using Abelian groups

A dipole lift construction of MMS graphs motivates the study of largest possible graphs of diameter 2 obtained as lifts of dipoles using voltages in Abelian groups.

In this case the upper bound can be improved as follows.

Lifts of dipoles using Abelian groups

A dipole lift construction of MMS graphs motivates the study of largest possible graphs of diameter 2 obtained as lifts of dipoles using voltages in Abelian groups.

In this case the upper bound can be improved as follows.

Theorem

[J. Šiagiová '02] For an arbitrary $d \geq 3$ let D be a dipole of degree d , and let α be a voltage assignment on D in an Abelian group such that the lift D^α is of diameter two. Then

$$|V(D^\alpha)| \leq \frac{4(10 + \sqrt{2})}{49}(d + 0.34)^2 \approx 0.932d^2$$

Lifting regular graphs

Lifting regular graphs

Let $K_n(m, l)$ denote a **complete graph** of order $n \geq 2$ with $l \geq 0$ loops on each vertex and $m \geq 1$ parallel edges between each pair of vertices.

Lifting regular graphs

Let $K_n(m, l)$ denote a **complete graph** of order $n \geq 2$ with $l \geq 0$ loops on each vertex and $m \geq 1$ parallel edges between each pair of vertices.

Lemma

Let α be an ordinary voltage assignment on a connected graph G in a group H and let k be a positive integer, and let u be a vertex of G . Then, the lift G^α has diameter at most k if and only if for every vertex v of G and for every element h of the group H there is a $u \rightarrow v$ path in G of length at most k such that $\alpha(W) = h$.

Lifting regular graphs

Let $K_n(m, l)$ denote a **complete graph** of order $n \geq 2$ with $l \geq 0$ loops on each vertex and $m \geq 1$ parallel edges between each pair of vertices.

Lemma

Let α be an ordinary voltage assignment on a connected graph G in a group H and let k be a positive integer, and let u be a vertex of G . Then, the lift G^α has diameter at most k if and only if for every vertex v of G and for every element h of the group H there is a $u \rightarrow v$ path in G of length at most k such that $\alpha(W) = h$.

Proposition

Let α be a voltage assignment on an n -pole $K_n(m, l)$ in an Abelian group A and suppose that the lift $K_n^\alpha(m, l)$ has diameter two. Then the number of vertices of $K_n^\alpha(m, l)$ is at most $\omega(m, l)$, where

$$\omega(m, l) = n \cdot \min\{1 + (n-1)m(m-1) + 2l(l+1), (n-2)m^2 + 4ml + m\}.$$

Theorem

Let $d \geq 3$ and $n \geq 2$ be integers and let α be a voltage assignment on an n -pole G of degree d in an Abelian group such that the lift G^α is of diameter 2. Then, for any fixed $n \geq 2$ the number of vertices of G^α is bounded above by

$$\frac{n^4 + 4n^3 + (2\sqrt{2} - 1)n^2 - (2\sqrt{2} + 2)n}{(n^2 + 2n - 1)^2} d^2 + O(d^{3/2})$$

as $d \rightarrow \infty$.

Sketch of proof:

We can express the polynomials from the Proposition in terms of degree d .


$$p_1(m) = \left(\frac{n^2}{2} - \frac{1}{2}\right)m^2 + (-dn - 2n + d + 2)m + \frac{d^2}{2} + d + 1, \quad \text{and}$$

$$p_2(m) = -nm^2 + (2d + 1)m, \quad \text{where in both cases } 1 \leq m \leq d/(n - 1).$$

Since we need to find

$$M = \max_m \min \{p_1(m), p_2(m)\}$$

we have two possibilities of intersection of the parabolas.



Considering the x -coordinates of the vertices of parabolas $p_1(m), p_2(m)$ we get that

$$M = \max_m \min \{p_1(m), p_2(m)\}$$

is attained at x_2 . Therefore

$$M = \max_m \min \{p_1(m), p_2(m)\} = p_1(x_2) = p_2(x_2)$$

If we substitute x_2 (expressed in terms of d, n) for m in $p_1(m)$ or $p_2(m)$ we gain the requesting result.

The upper bound for a diameter-two lift of $K_n(m, l)$ for some values of n :

n	Order of lift of n -pole
2	$0.932d^2 + O(d)$
3	$0.974d^2 + O(d)$
4	$0.987d^2 + O(d)$
5	$0.992d^2 + O(d)$
6	$0.995d^2 + O(d)$
7	$0.996d^2 + O(d)$

Conclusion

Our aim now is to investigate diameter-two lifts of other regular graphs on n vertices. We wish either to show that the upper bound obtained from the lifts of $K_n(m, l)$ is the best possible or to find a new regular graph on n vertices the lift of which has bigger order than presented.

THANK YOU FOR YOUR ATTENTION