

Internal Operators: Application to Image Processing

D. Paternain, J. Fernandez, H. Bustince

Public University of Navarra

Pamplona, Spain

G. Beliakov

Deakin University

Burwood, Australia

May 12, 2012

Outline

Motivation

Preliminaries

Internal operators. Definition and properties

Construction of internal operators from penalty functions

Conclusions

Motivation

Image Processing:

Motivation

Image Processing:

- ▶ The need to aggregate several values into one representative is very important in several applications:

Motivation

Image Processing:

- ▶ The need to aggregate several values into one representative is very important in several applications:
- ▶ Image filtering

Motivation

Image Processing:

- ▶ The need to aggregate several values into one representative is very important in several applications:
- ▶ Image filtering
- ▶ Stereo vision

Motivation

Image Processing:

- ▶ The need to aggregate several values into one representative is very important in several applications:
- ▶ Image filtering
- ▶ Stereo vision
- ▶ Image reduction-reconstruction

Motivation

Image Processing:

- ▶ The need to aggregate several values into one representative is very important in several applications:
- ▶ Image filtering
- ▶ Stereo vision
- ▶ Image reduction-reconstruction
- ▶ Which is the best aggregated value?

Motivation


Image Processing:

- ▶ The need to aggregate several values into one representative is very important in several applications:
- ▶ Image filtering
- ▶ Stereo vision
- ▶ Image reduction-reconstruction
- ▶ Which is the best aggregated value?

Example: Image Reduction









Preliminaries

Definition

A function $M : [0, 1]^n \rightarrow [0, 1]$ is called an n -ary aggregation function if it is monotone non-decreasing in each variable and such that $M(0, \dots, 0) = 0$ and $M(1, \dots, 1) = 1$.

Definition

We say that an aggregation function M is

- ▶ idempotent if $M(x, \dots, x) = x$ for all $x \in [0, 1]$.
- ▶ averaging if $\min(x_1, \dots, x_n) \leq M(x_1, \dots, x_n) \leq \max(x_1, \dots, x_n)$ for every $(x_1, \dots, x_n) \in [0, 1]^n$.
- ▶ conjunctive if $M(x_1, \dots, x_n) \leq \min(x_1, \dots, x_n)$.
- ▶ disjunctive if $M(x_1, \dots, x_n) \geq \max(x_1, \dots, x_n)$.
- ▶ mixed if it is not conjunctive, disjunctive nor averaging, that is, it has different behaviours on different parts of the domain.
- ▶ symmetric if $M(x_1, \dots, x_n) = M(x_{\sigma(1)}, \dots, x_{\sigma(n)})$ for every $(x_1, \dots, x_n) \in [0, 1]^n$ and every permutation $\sigma = (\sigma(1), \dots, \sigma(n))$.

Some families:

Definition

Let $g : [0, 1] \rightarrow [-\infty, \infty]$ be a continuous and strictly monotone function. We call quasi-arithmetic mean the mapping

$M_g : [0, 1]^n \rightarrow [0, 1]$ defined as

$$M_g(x_1, \dots, x_n) = g^{-1} \left(\frac{1}{n} \sum_{i=1}^n g(x_i) \right)$$

Some families:

Definition

Let $g : [0, 1] \rightarrow [-\infty, \infty]$ be a continuous and strictly monotone function. We call quasi-arithmetic mean the mapping

$M_g : [0, 1]^n \rightarrow [0, 1]$ defined as

$$M_g(x_1, \dots, x_n) = g^{-1} \left(\frac{1}{n} \sum_{i=1}^n g(x_i) \right)$$

Definition

A mapping $F : [0, 1]^n \rightarrow [0, 1]$ is called an OWA operator of dimension n if there exists a weighting vector

$W = (w_1, \dots, w_n) \in [0, 1]^n$ with $\sum_i w_i = 1$ and such that

$$F(x_1, \dots, x_n) = \sum_{i=1}^n w_i x_{(i)}$$

Internal operators. Definition and properties

Definition

An internal operator is a mapping $F : [0, 1]^n \rightarrow [0, 1]$ such that

$$F(x_1, \dots, x_n) \in \{x_1, \dots, x_n\}.$$

1. An internal operator needs not to be homogeneous. Consider the following IF operator:

$$IF(x_1, \dots, x_n) = \begin{cases} \min(x_1, \dots, x_n) & \text{if } \max(x_1, \dots, x_n) \leq \frac{1}{4} \\ x_1 & \text{otherwise} \end{cases}$$

Then, if we take $n = 3$ and $\lambda = 0.5$, we have that

$$IF\left(\frac{1}{2}, \frac{1}{3}, \frac{1}{4}\right) = \frac{1}{4} \text{ and } IF\left(\lambda\frac{1}{2}, \lambda\frac{1}{3}, \lambda\frac{1}{4}\right) = \frac{1}{4} \neq \lambda IF\left(\frac{1}{2}, \frac{1}{3}, \frac{1}{4}\right).$$

1. An internal operator needs not to be homogeneous. Consider the following IF operator:

$$IF(x_1, \dots, x_n) = \begin{cases} \min(x_1, \dots, x_n) & \text{if } \max(x_1, \dots, x_n) \leq \frac{1}{4} \\ x_1 & \text{otherwise} \end{cases}$$

Then, if we take $n = 3$ and $\lambda = 0.5$, we have that

$$IF\left(\frac{1}{2}, \frac{1}{3}, \frac{1}{4}\right) = \frac{1}{4} \text{ and } IF\left(\lambda\frac{1}{2}, \lambda\frac{1}{3}, \lambda\frac{1}{4}\right) = \frac{1}{4} \neq \lambda IF\left(\frac{1}{2}, \frac{1}{3}, \frac{1}{4}\right).$$

2. An internal operator needs not to be shift-invariant. Consider the same example as below. Then for $n = 3$ and $r = \frac{1}{2}$ we have that $IF\left(\frac{1}{4}, \frac{1}{6}, \frac{1}{8}\right) = \frac{1}{8}$ and $IF\left(\frac{1}{2} + r, \frac{1}{3} + r, \frac{1}{4} + r\right) = \frac{3}{4} \neq IF\left(\frac{1}{2}, \frac{1}{3}, \frac{1}{4}\right) + r.$

1. An internal operator needs not to be homogeneous. Consider the following IF operator:

$$IF(x_1, \dots, x_n) = \begin{cases} \min(x_1, \dots, x_n) & \text{if } \max(x_1, \dots, x_n) \leq \frac{1}{4} \\ x_1 & \text{otherwise} \end{cases}$$

Then, if we take $n = 3$ and $\lambda = 0.5$, we have that

$$IF\left(\frac{1}{2}, \frac{1}{3}, \frac{1}{4}\right) = \frac{1}{4} \text{ and } IF\left(\lambda\frac{1}{2}, \lambda\frac{1}{3}, \lambda\frac{1}{4}\right) = \frac{1}{4} \neq \lambda IF\left(\frac{1}{2}, \frac{1}{3}, \frac{1}{4}\right).$$

2. An internal operator needs not to be shift-invariant. Consider the same example as below. Then for $n = 3$ and $r = \frac{1}{2}$ we have that $IF\left(\frac{1}{4}, \frac{1}{6}, \frac{1}{8}\right) = \frac{1}{8}$ and $IF\left(\frac{1}{2} + r, \frac{1}{3} + r, \frac{1}{4} + r\right) = \frac{3}{4} \neq IF\left(\frac{1}{2}, \frac{1}{3}, \frac{1}{4}\right) + r$.
3. An internal operator needs not to be monotone. When there are at least one input repeated, the mode is an example of a non-monotone internal operator.

Proposition

Let F be an internal operator. The following items hold:

- i) F is idempotent;
- ii) F is compensative, that is,

$$\min(x_1, \dots, x_n) \leq F(x_1, \dots, x_n) \leq \max(x_1, \dots, x_n).$$

Internal Aggregation Functions

Theorem

Let $F : [0, 1]^n \rightarrow [0, 1]$ be an internal operator. If F is monotone non-decreasing in each variable, then F is an averaging aggregation function.

Internal Aggregation Functions

Theorem

Let $F : [0, 1]^n \rightarrow [0, 1]$ be an internal operator. If F is monotone non-decreasing in each variable, then F is an averaging aggregation function.

Corollary

Under conditions of Theorem 6, F is jointly strict monotone, that is $x_i < y_i$ for all $i \in \{1, \dots, n\}$ implies $F(x_1, \dots, x_n) < F(y_1, \dots, y_n)$.

Theorem

The following items hold:

- i) *There is no conjunctive internal operators other than the minimum.*
- ii) *There is no disjunctive internal operators other than the maximum.*

Theorem

Let $M_g : [0, 1]^n \rightarrow [0, 1]$ be a quasi-arithmetic mean. The following items are equivalent:

- i) M_g is internal operator;
- ii) there exist $x_j \in \{x_1, \dots, x_n\}$ such that

$$x_j = M_g(x_1, \dots, x_{j-1}, x_{j+1}, \dots, x_n) = g^{-1} \left(\frac{1}{n-1} \sum_{\substack{i=1 \\ i \neq j}}^n g(x_i) \right).$$

Theorem

Let $M_g : [0, 1]^n \rightarrow [0, 1]$ be a quasi-arithmetic mean. The following items are equivalent:

- i) M_g is internal operator;
- ii) there exist $x_j \in \{x_1, \dots, x_n\}$ such that

$$x_j = M_g(x_1, \dots, x_{j-1}, x_{j+1}, \dots, x_n) = g^{-1} \left(\frac{1}{n-1} \sum_{\substack{i=1 \\ i \neq j}}^n g(x_i) \right).$$

Corollary

There is not any internal quasi-arithmetic mean for all $(x_1, \dots, x_n) \in [0, 1]^n$.

Theorem

Let $F : [0, 1]^n \rightarrow [0, 1]$ be an OWA operator of dimension n such that $w_j \in [0, 1[$ for all $j \in \{1, \dots, n\}$. F is internal if and only if there exists $x_{(j)} \in \{x_{(1)}, \dots, x_{(n)}\}$ such that

$$x_{(j)} = \sum_{\substack{i=1 \\ i \neq j}}^n \frac{w_i}{1 - w_j} x_{(i)}.$$

Theorem

Let $F : [0, 1]^n \rightarrow [0, 1]$ be an OWA operator of dimension n such that $w_j \in [0, 1[$ for all $j \in \{1, \dots, n\}$. F is internal if and only if there exists $x_{(j)} \in \{x_{(1)}, \dots, x_{(n)}\}$ such that

$$x_{(j)} = \sum_{\substack{i=1 \\ i \neq j}}^n \frac{w_i}{1 - w_j} x_{(i)}.$$

Corollary

Let $F : [0, 1]^n \rightarrow [0, 1]$ be an OWA operator of dimension n associated with the weighting vector $w = (w_1, \dots, w_n)$. F is internal if and only if there exists $i \in \{1, \dots, n\}$ such that $w_i = 1$ and $w_j = 0$ for every $j \neq i$.

Internal operators and penalty functions

In a recent paper, we have studied the aggregation of color pixels by means of Penalty Functions.

How can we define aggregation functions by means of penalty functions?

- ▶ The arithmetic mean and the median are the solution to optimization problems;

Internal operators and penalty functions

In a recent paper, we have studied the aggregation of color pixels by means of Penalty Functions.

How can we define aggregation functions by means of penalty functions?

- ▶ The arithmetic mean and the median are the solution to optimization problems;
- ▶ Suppose we have the input $\mathbf{x} = (x_1, \dots, x_n)$ and a possible output y ;

Internal operators and penalty functions

In a recent paper, we have studied the aggregation of color pixels by means of Penalty Functions.

How can we define aggregation functions by means of penalty functions?

- ▶ The arithmetic mean and the median are the solution to optimization problems;
- ▶ Suppose we have the input $\mathbf{x} = (x_1, \dots, x_n)$ and a possible output y ;
- ▶ We want to measure the disagreement between input and output;

Internal operators and penalty functions

In a recent paper, we have studied the aggregation of color pixels by means of Penalty Functions.

How can we define aggregation functions by means of penalty functions?

- ▶ The arithmetic mean and the median are the solution to optimization problems;
- ▶ Suppose we have the input $\mathbf{x} = (x_1, \dots, x_n)$ and a possible output y ;
- ▶ We want to measure the disagreement between input and output;
- ▶ If some input $x_i \neq y$, we impose a penalty for the disagreement.

Definition

Let $P : [a, b]^{n+1} \rightarrow \mathcal{R}$ be a penalty function with the properties

- i) $P(\mathbf{x}, y) \geq 0$ for all \mathbf{x}, y ;
- ii) $P(\mathbf{x}, y) = 0$ if all $x_i = y$;
- iii) $P(x, y)$ is quasi-convex in y for any \mathbf{x} .

Definition

Let $P : [a, b]^{n+1} \rightarrow \mathcal{R}$ be a penalty function with the properties

- i) $P(\mathbf{x}, y) \geq 0$ for all \mathbf{x}, y ;
- ii) $P(\mathbf{x}, y) = 0$ if all $x_i = y$;
- iii) $P(\mathbf{x}, y)$ is quasi-convex in y for any \mathbf{x} .

The penalty based function is

$$f(\mathbf{x}) = \arg \min_y P(\mathbf{x}, y)$$

if y is the unique minimizer, and $y = \frac{a+b}{2}$ if the set of minimizers is the interval $[a, b]$.

Definition

Let $P : [a, b]^{n+1} \rightarrow \mathcal{R}$ be a penalty function with the properties

- i) $P(\mathbf{x}, y) \geq 0$ for all \mathbf{x}, y ;
- ii) $P(\mathbf{x}, y) = 0$ if all $x_i = y$;
- iii) $P(\mathbf{x}, y)$ is quasi-convex in y for any \mathbf{x} .

The penalty based function is

$$f(\mathbf{x}) = \arg \min_y P(\mathbf{x}, y)$$

if y is the unique minimizer, and $y = \frac{a+b}{2}$ if the set of minimizers is the interval $[a, b]$.

Theorem

Any averaging aggregation function can be expressed as a penalty based function.

Examples:

- ▶ The arithmetic mean is the value that minimizes

$$f(\mathbf{x}) = \arg \min_y \sum_{i=1}^n (x_i - y)^2$$

Examples:

- ▶ The arithmetic mean is the value that minimizes

$$f(\mathbf{x}) = \arg \min_y \sum_{i=1}^n (x_i - y)^2$$

- ▶ The median is the value that minimizes

$$f(\mathbf{x}) = \arg \min_y \sum_{i=1}^n |x_i - y|$$

Definition

Let $P : [0, 1]^{n+1} \rightarrow [0, 1]$ be a penalty function. We call internal penalty based function F to

$$F(x_1, \dots, x_n) = \arg \min_{x_i} P(\mathbf{x}, x_i)$$

Theorem

Let $F : [0, 1]^n \rightarrow [0, 1]$ be an internal penalty based function given by

$$F(x_1, \dots, x_n) = \arg \min_{x_i} P(\mathbf{x}, x_i).$$

Then, F is an internal operator.

We are using this construction method to build internal aggregation functions:

We are using this construction method to build internal aggregation functions:

- ▶ The internal version of the arithmetic mean:

$$f(\mathbf{x}) = \arg \min_{x_j} \sum_{i=1}^n (x_i - x_j)^2$$

We are using this construction method to build internal aggregation functions:

- ▶ The internal version of the arithmetic mean:

$$f(\mathbf{x}) = \arg \min_{x_j} \sum_{i=1}^n (x_i - x_j)^2$$

- ▶ The internal version of the median

$$f(\mathbf{x}) = \arg \min_{x_j} \sum_{i=1}^n |x_i - x_j|$$

We are using this construction method to build internal aggregation functions:

- ▶ The internal version of the arithmetic mean:

$$f(\mathbf{x}) = \arg \min_{x_j} \sum_{i=1}^n (x_i - x_j)^2$$

- ▶ The internal version of the median

$$f(\mathbf{x}) = \arg \min_{x_j} \sum_{i=1}^n |x_i - x_j|$$

- ▶ The internal version of the geometric mean

$$f(\mathbf{x}) = \arg \min_{x_j} \sum_{i=1}^n (-\log(x_i) + \log(x_j))^2$$

Conclusions

- ▶ We have seen the need of using internal aggregation functions in some processes of image processing
- ▶ We have studied the conditions under which well known aggregation functions are internal or not
- ▶ This study has led us to study a way of constructing new internal aggregation functions
- ▶ We have given a way of constructing these functions by minimizing special penalty functions